

Dietary Reference Intakes (DRIs): Recommended Dietary Allowances and Adequate Intakes, Total Water and Macronutrients

Food and Nutrition Board, Institute of Medicine, National Academies

Life Stage Group	Total Water ^a (L/d)	Carbohydrate (g/d)	Total Fiber (g/d)	Fat (g/d)	Linoleic Acid (g/d)	α -Linolenic Acid (g/d)	Protein ^b (g/d)
Infants							
0–6 mo	0.7*	60*	ND	31*	4.4*	0.5*	9.1*
6–12 mo	0.8*	95*	ND	30*	4.6*	0.5*	11.0
Children							
1–3 y	1.3*	130	19*	ND ^c	7*	0.7*	13
4–8 y	1.7*	130	25*	ND	10*	0.9*	19
Males							
9–13 y	2.4*	130	31*	ND	12*	1.2*	34
14–18 y	3.3*	130	38*	ND	16*	1.6*	52
19–30 y	3.7*	130	38*	ND	17*	1.6*	56
31–50 y	3.7*	130	38*	ND	17*	1.6*	56
51–70 y	3.7*	130	30*	ND	14*	1.6*	56
> 70 y	3.7*	130	30*	ND	14*	1.6*	56
Females							
9–13 y	2.1*	130	26*	ND	10*	1.0*	34
14–18 y	2.3*	130	26*	ND	11*	1.1*	46
19–30 y	2.7*	130	25*	ND	12*	1.1*	46
31–50 y	2.7*	130	25*	ND	12*	1.1*	46
51–70 y	2.7*	130	21*	ND	11*	1.1*	46
> 70 y	2.7*	130	21*	ND	11*	1.1*	46
Pregnancy							
14–18 y	3.0*	175	28*	ND	13*	1.4*	71
19–30 y	3.0*	175	28*	ND	13*	1.4*	71
31–50 y	3.0*	175	28*	ND	13*	1.4*	71
Lactation							
14–18	3.8*	210	29*	ND	13*	1.3*	71
19–30 y	3.8*	210	29*	ND	13*	1.3*	71
31–50 y	3.8*	210	29*	ND	13*	1.3*	71


NOTE: This table (take from the [DRI reports](http://www.nap.edu), see www.nap.edu) presents Recommended Dietary Allowances (RDA) in **bold type** and Adequate Intakes (AI) in ordinary type followed by an asterisk (*). An RDA is the average daily dietary intake level sufficient to meet the nutrient requirements of nearly all (97–98 percent) healthy individuals in a group. It is calculated from an Estimated Average Requirement (EAR). If sufficient scientific evidence is not available to establish an EAR, and thus calculate an RDA, an AI is usually developed. For healthy breast-fed infants, an AI is the mean intake. The AI for other life stage and gender groups is believed to cover the needs of all

healthy individuals in the groups, but lack of data or uncertainty in the data prevent being able to specify with confidence the percentage of individuals covered by this intake.

- a Total water includes all water contained in food, beverages, and drinking water.
- b Based on g protein per kg of body weight for the reference body weight, e.g., for adults 0.8 g/kg body weight for the reference body weight.
- c Not determined.

SOURCE: *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids* (2002/2005) and *Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate* (2005). The report may be accessed via www.nap.edu.

From: Summary Tables



Dietary Reference Intakes for Calcium and Vitamin D.
Institute of Medicine (US) Committee to Review Dietary Reference Intakes for Vitamin D and Calcium; Ross AC, Taylor CL, Yaktine AL, et al., editors.
Washington (DC): National Academies Press (US); 2011.

Copyright © 2011, National Academy of Sciences.

NCBI Bookshelf. A service of the National Library of Medicine, National Institutes of Health.